

Lecture Four

(代数计算)

本节课授课要点

代数计算

- 指数运算
- 解方程
- 不等式
- 符号运算
- 数列

指数运算 (Rules of Exponents)

$$a^m \times a^n = a^{m+n}$$

$$a^m \div a^n = a^{m-n}$$

$$(a^m)^n = a^{mn}$$

$$a^m \times b^m = (a \times b)^m$$

$$a^m \div b^m = (a \div b)^m$$

1. $\frac{(8)^2(3)^3(2)^4}{(96)^2} = ?$

(A) 3

(B) 6

(C) 9

(D) 12

(E) 18

2. If $P = 2^{64}$ and $P^P = 2^K$, then $k =$

- (A) 70
- (B) 128
- (C) 256
- (D) 2^{64}
- (E) 2^{70}

3. The function f is defined for each positive three-digit integer n by $f(n) = 2^x 3^y 5^z$, where x , y and z are the hundreds, tens, and units digits of n , respectively. If m and v are three-digit positive integers such that $f(m) = 9f(v)$, then $m - v = ?$

- (A) 8
- (B) 9
- (C) 18
- (D) 20
- (E) 80

本节课授课要点

代数计算

- 指数运算
- 解方程
- 不等式
- 符号运算
- 数列

解方程 (Equations)

一元二次方程 $ax^2 + bx + c = 0$

标准根的公式为: $x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

1. A square playground has the same area as a rectangular playground that is 30 meters longer but 20 meters narrower. What is the length, in meters, of a side of the square playground?

- (A) $10\sqrt{5}$
- (B) $10\sqrt{6}$
- (C) 25
- (D) 50
- (E) 60

2. A certain theater has 100 balcony seats. For every \$2 increase in the price of a balcony seat above \$10, 5 fewer seats will be sold. If all the balcony seats are sold when the price of each seat is \$10, which of the following could be the price of a balcony seat if the revenue from the sale of balcony seats is \$1,360 ?

- (A) \$12
- (B) \$14
- (C) \$16
- (D) \$17
- (E) \$18

3. If the sum of two positive integers is 24 and the difference of their squares is 48, what is the product of the two integers?

- (A) 108
- (B) 119
- (C) 128
- (D) 135
- (E) 143

4. If $x - \sqrt{10} = \sqrt{x} + 10$, then $x =$

5. What is Steve's annual salary and Maria's annual salary?

(1) The combined total of the annual salaries of Steve and Maria is \$80,000.

(2) If Steve were to receive a 10 percent increase in annual salary and Maria a 10 percent increase, their combined annual salaries would be \$88,000.

本节课授课要点

代数计算

- 指数运算
- 解方程
- 不等式
- 符号运算
- 数列

对已有的不等式两边取倒数或负数，不等号通常要改变方向

对 $\sqrt[4]{x}, \sqrt[3]{x}, \sqrt{x}, x, x^2, x^3, x^4$ 等函数的性质有一定的认识.

在 x, x^2, x^3 几个函数的比较大小中, 对 x 的取值范围要有清醒的分段意识:

$$x < -1, \quad -1 < x < 0, \quad 0 < x < 1, \quad x > 1$$

绝对值: $|x|$ 恒非负

1. If $x > 0.9$, which of the following could be the value of x ?

(A) $\sqrt{0.81}$

(B) $\sqrt{0.9}$

(C) (0.9)

(D) $(0.9)(0.9)$

(E) $1 - \sqrt{0.01}$

2. If $x \neq 0$, is $|x| < 1$?

(1) $x^2 < 1$

(2) $|x| < \frac{1}{x}$

3. Which of the following inequalities has a solution set that, when graphed in the number line, is a single line segment of finite length?

(A) $x^4 \geq 16$

(B) $x^3 \leq 27$

(C) $x^2 \geq 16$

(D) $2 \leq |x| \leq 5$

(E) $2 \leq 3x + 4 \leq 6$

本节课授课要点

代数计算

- 指数运算
- 解方程
- 不等式
- 符号运算
- 数列

题目会定义新的算符和运算法则，模仿运算

1. For all real numbers v , the operation v^* is defined by the equation $v^* = v - v/3$. If $(v^*)^* = 8$, then $v =$

- (A) 15
- (B) 18
- (C) 21
- (D) 24
- (E) 27

2. If the operation \odot is defined for all integers a and b by $a \odot b = a + b - ab$, which of the following statements must be true for all integers a , b , and c ?

I. $a \odot b = b \odot a$

II. $a \odot 0 = a$

III. $(a \odot b) \odot c = a \odot (b \odot c)$

(A) I only

(B) II only

(C) I and II only

(D) I and III only

(E) I, II and III

3. If \odot denotes one of two arithmetic operations, addition or multiplication, and if k is an integer, what is the value of $3 \odot k$?

(1) $2 \odot k = 3$

(2) $1 \odot 0 = k$

本节课授课要点

代数计算

- 指数运算
- 解方程
- 不等式
- 符号运算
- 数列

1. 等差数列 (Arithmetic Sequence)

2. 等比数列 (Geometric Sequence)

1. How many integers between 234 and 567, inclusive, are the multiples of 5 ?

- (A) 62
- (B) 65
- (C) 67
- (D) 69
- (E) 71

2. If the sum of 7 consecutive integers is 434, then the greatest of the 7 integers is

- (A) 65
- (B) 66
- (C) 67
- (D) 68
- (E) 69

3. In the sequence 1, 2, 4, 8, 16, 32, ... , each term after the first is twice the previous term. What is the sum of the 16th, 17th, and 18th terms in the sequence?

- (A) 2^{18}
- (B) $3(2^{17})$
- (C) $7(2^{16})$
- (D) $3(2^{16})$
- (E) $7(2^{15})$

4. In a certain sequence, the first term is 1, and each successive term is 1 more than the reciprocal of the term that immediately precedes it. What is the fifth term of the sequence?

- (A) $\frac{3}{5}$ (B) $\frac{5}{8}$ (C) $\frac{8}{5}$ (D) $\frac{5}{3}$ (E) $\frac{9}{2}$

5. The sequence $a_1, a_2, \dots, a_n, \dots$ is such that $a_n = a_{n-1} - a_{n-2}$ for all positive integers $n > 2$. If $a_1 = -1$ and $a_2 = 1$, what is the sum of the first 1000 terms in the sequence?

- (A) 0
- (B) 3
- (C) 750
- (D) 1000
- (E) 3000

回顾本节课授课要点

代数计算

- 指数运算
- 解方程
- 不等式
- 符号运算
- 数列

预告下节课授课要点

初等几何

- 三角形与四边形
- 平行线
- 圆
- 立体几何
- 直角坐标系

The End